

K -causality coincides with stable causality

E. Minguzzi

Dipartimento di Matematica Applicata, Università degli Studi di Firenze, Via S. Marta 3,
I-50139 Firenze, Italy
E-mail: ettore.minguzzi@unifi.it

Abstract: It is proven that K -causality coincides with stable causality, and that in a K -causal spacetime the relation K^+ coincides with the Seifert's relation. As a consequence the causal relation “the spacetime is strongly causal and the closure of the causal relation is transitive” stays between stable causality and causal continuity.

1. Introduction

The relation K^+ is defined as the smallest closed and transitive relation which contains the causal relation J^+ . It was introduced by Sorkin and Woolgar in [16] who also defined a spacetime as K -causal if the relation K^+ is antisymmetric.

The relation K^+ was originally conceived to recast global causal analysis in an order-theoretic framework, or as a tool for exploring spacetimes with C^0 metrics or varying topology [16,3]. In [7,8] I compared K -causality with the levels already present in the causal ladder of spacetimes [15,13], a well known hierarchy of conformal invariant properties whose study started in a seminal work by Hawking and Sachs [5], and which in the last years has seen the introduction of new levels [7,11] and some improvements [13,2,12].

In this respect since the introduction of K -causality R. Low [16, footnote p. 1990] suggested the coincidence of this relation with stable causality [4]. Indeed, stable causality is equivalent to the antisymmetry of Seifert's relation [14] $J_S^+ = \bigcap_{g' > g} J_{g'}^+$ (a fact rigorously proved in [8], see also [5]) where J_S^+ is a closed, transitive relation which contains J^+ . These last properties imply, from the definition of K^+ , $K^+ \subset J_S^+$, and since the antisymmetric property is inherited by inclusion, stable causality implies K -causality.

The open question was whether the equality $K^+ = J_S^+$ holds, because in this case K -causality and stable causality would be equivalent. Actually [8] there are

causal examples for which $J_S^+ \neq K^+$, but nevertheless it could still be that K -causality coincides with stable causality, in particular if K -causality forces the equivalence $K^+ = J_S^+$. In Seifert's work [14] there is indeed an unproved claim¹ (lemma 2) which is equivalent to such a statement, although it should be noted that K -causality was not yet defined at the time (see [8] for a discussion). Thus the problem of the equivalence between stable and K -causality has been around for almost four decades, though it has attracted attention only in the last twelve years.

As I shall prove below, K -causality and stable causality do indeed coincide and, thanks to the results of [8], this equivalence implies that in a K -causal spacetime the K^+ relation coincides with the Seifert relation. Given this result the logical structure of some other proofs simplify considerably, I mention the proof that causal continuity implies stable causality and the proof that chronological spacetimes without lightlike lines are stably causal. It also suggests the definition of a new causal relation which stays between stable causality and causal continuity. This relation, here termed *causal easiness*, is: the spacetime is strongly causal and $\bar{J}^+$ is transitive.

The proof of the coincidence between stable causality and K -causality uses the concept of “compact stable causality” introduced in [9]. In short a spacetime is compactly stably causal if for every compact set the light cones can be widened on the compact set while preserving causality. In [9] I proved that K -causality implies compact stable causality, and I gave examples which show that the two properties differ.

I refer the reader to [13, 7] for most of the conventions used in this work. In particular, I denote with (M, g) a C^r spacetime (connected, time-oriented Lorentzian manifold), $r \in \{3, \dots, \infty\}$ of arbitrary dimension $n \geq 2$ and signature $(-, +, \dots, +)$. On $M \times M$ the usual product topology is defined. For convenience and generality I often use the causal relations on $M \times M$ in place of the more widespread point based relations $I^+(x)$, $J^+(x)$, $E^+(x)$ (and past versions). All the causal curves that we shall consider are future directed. The subset symbol $\subset$ is reflexive, $X \subset X$. Several versions of the limit curve theorem will be repeatedly used, particularly those referring to sequences of g_n -causal curves, where the metrics in the sequence g_n may differ. The reader is referred to [10] for a sufficiently strong formulation. With A^+ I denote [17, 1, 7] the closure of the causal relation, $A^+ = \bar{J}^+$, and a spacetime on which A^+ is antisymmetric is called A -causal. For our purposes, it will be useful to recall the implications: K -causality $\Rightarrow$ compact stable causality $\Rightarrow A$ -causality $\Rightarrow$ strong causality $\Rightarrow$ non-total imprisonment $\Rightarrow$ causality. Subsequences are denoted by changing the index, thus x_k may denote a subsequence of x_n . The set Δ is the diagonal on $M \times M$.

¹ Seifert's unproved claim has raised some confusion in recent literature. I warn the reader that in the preprint gr-qc/9912090v1, Dowker et al. claimed that stable causality implies $K^+ = J_S^+$. Actually, the proof relied on the Seifert's lemma 2, so that after realizing the inconsistency of that lemma they correctly removed this statement from the published version [3]. Unfortunately, in [6] the authors attribute this result to Dowker et al., as they took this information from the preprint version.

2. K -causality coincides with stable causality

We need some preliminary lemmas. The first one basically states that if two points are K -related but not causally related then it is possible to find a new point, in a compact shell as close to infinity as one wishes, which stays in the “middle” of the original points.

Lemma 1. *Let (M, g) be a non-total imprisoning spacetime. If $(x, z) \in K^+ \setminus J^+$ then for every compact C there is $w \in M \setminus C$ such that $(x, w) \in K^+$ and $(w, z) \in K^+$. In particular if $(x, z) \in K^+ \setminus J^+$ then for every open set with compact closure B , with $x, z \in B$, there is $y \in \dot{B}$ such that $(x, y) \in K^+$ and $(y, z) \in K^+$.*

Proof. Let

$$R^+ = \{(x, z) \in K^+ \text{ such that } (x, z) \in J^+ \text{ or for every compact set } C \\ \text{there is } w \in M \setminus C \text{ such that } (x, w) \in K^+ \text{ and } (w, z) \in K^+\}$$

we are going to prove that R^+ is closed and transitive, and since $J^+ \subset R^+ \subset K^+$ this fact will imply $R^+ = K^+$ from which the first statement will follow. The last statement is a trivial consequence of the first statement and [16, lemma 14] [8, lemma 5.3].

For the transitivity let $(x, y) \in R^+$ and $(y, z) \in R^+$. If both belong to J^+ then $(x, z) \in J^+ \subset R^+$. If the latter pair does not belong to J^+ then whatever the compact set C there is $w \in M \setminus C$ such that $(y, w) \in K^+$ and $(w, z) \in K^+$ thus since $(x, y) \in K^+$, we have $(x, w) \in K^+$ and hence $(x, z) \in R^+$. If the former pair does not belong to J^+ the proof is analogous.

For the closure let $(x_n, z_n) \rightarrow (x, z)$ with $(x_n, z_n) \in R^+$. We have to prove that $(x, z) \in R^+$ thus we can assume $x \neq z$, since $\Delta \subset J^+ \subset R^+$. If there is a subsequence $(x_k, z_k) \in J^+$ let σ_k be a sequence of causal curves connecting x_k to z_k . By the limit curve theorem either there is a causal curve connecting x to z , in which case $(x, z) \in J^+ \subset R^+$, and there is nothing left to prove, or there is a past inextendible limit causal curve σ^z ending at z , such that for every point $w \in \sigma^z$, $(x, w) \in J^+ \subset K^+$. Since (M, g) is non-total imprisoning σ^z must escape every compact, thus chosen a compact C , w can be chosen in $M \setminus C$. Since clearly $(w, z) \in J^+ \subset K^+$ it follows $(x, z) \in R^+$.

Thus without loss of generality we can assume that none of the elements in the sequence (x_n, z_n) belong to J^+ . Let C be a compact and let B be a open set with compact closure such that $C \subset B$ and $x, z \in B$ so that we can assume (pass to a subsequence if necessary) $x_n, z_n \in B$. Since $(x_n, z_n) \in R^+$ and $\bar{B}$ is compact, there is w'_n in $M \setminus \bar{B}$ such that $(x_n, w'_n) \in K^+$ and $(w'_n, z_n) \in K^+$. By a well known result [16, lemma 14] [8, lemma 5.3] it is possible to find $w_n \in \dot{B} \subset M \setminus C$ such that $(x_n, w_n) \in K^+$ and $(w_n, z_n) \in K^+$. Since $\dot{B}$ is compact there is a subsequence such that $(x_i, w_i) \rightarrow (x, w)$ and $(w_i, z_i) \rightarrow (w, z)$ with $w \in \dot{B}$. Since K^+ is closed we have in particular $(x, w) \in K^+$ and $(w, z) \in K^+$ from which $(x, z) \in R^+$ follows.

The next lemma clarifies that if it is possible to enlarge the light cones in an arbitrary compact set while preserving K -causality then the process can be continued all over the spacetime.

Lemma 2. *The statement “if (M, g) is K -causal then for every compact set C , there is a metric $g_C \geq g$, such that $g_C > g$ on C , with (M, g_C) K -causal”, implies the apparently stronger statement “if (M, g) is K -causal then it is also stably causal”.*

Proof. Assume the first statement. Note that if $C \subset \text{Int } C'$, and C' is compact, by taking a point dependent convex combination of g and g_C it is possible to find $g'_C \geq g$, $g < g'_C \leq g_C$ on C , $g'_C = g$ outside C' . Hence, since $g'_C \leq g_C$, (M, g'_C) is K -causal (recall lemma 5.10 of [8]). Thus the statement “if (M, g) is K -causal then for every compact C , there is a metric $g_C \geq g$, such that $g_C > g$ on C , with (M, g_C) K -causal” implies “if (M, g) is K -causal then for every pair of compacts C, C' , $C \subset \text{Int } C'$, there is a metric $g_{CC'} \geq g$, such that $g_{CC'} > g$ on C , $g_{CC'} = g$ outside C' with $(M, g_{CC'})$ K -causal”

Assume that (M, g) is K -causal. Take $p \in M$ and let h be a complete Riemannian metric on M . Let $B_n(p)$ be closed balls centered at p of h -radius n . By the Hopf-Rinow theorem they are compact. Let $g_2 \geq g$ be a metric such that $g_2 > g$ on $B_2(p)$, $g_2 = g$ outside $B_3(p)$ and (M, g_2) is K -causal. Consider the compacts $C_3 = B_3(p) \setminus \text{Int } B_2(p)$, and $C'_3 = B_4(p) \setminus \text{Int } B_1(p)$ let $g_3 \geq g_2$ be a metric such that $g_3 > g_2$ on C_3 , $g_3 = g_2$ outside C'_3 , and (M, g_3) is K -causal. Continue in this way by defining $C_n = B_n(p) \setminus \text{Int } B_{n-1}(p)$, and $C'_n = B_{n+1}(p) \setminus \text{Int } B_{n-2}(p)$, and let $g_n \geq g_{n-1}$ be a metric such that $g_n > g_{n-1}$ on C_n , $g_n = g_{n-1}$ outside C'_n . By induction given the assumed statement, (M, g_n) is K -causal. Now, note that if $x \in B_n(p)$ then $g_k(x)$ is independent of k for $k \geq n+1$. Define g' so that if $x \in B_n(p)$, $g'(x) = g_{n+1}(x)$. Clearly, for every n , $g' \geq g_n$ and $g' > g$. Suppose (M, g') is not causal then there is a closed g' -causal curve γ , which necessarily is contained in a compact $B_s(p)$. But $g'|_{B_s(p)} = g_{s+1}|_{B_s(p)}$, thus γ is g_{s+1} -causal in contradiction with the causality of (M, g_{s+1}) . Thus (M, g) is stably causal.

In order to prove that the metric can be enlarged over a compact set C while preserving K -causality, we are going to enlarge it in a finite covering of C made of open sets A_x constructed as in the next lemma.

As a matter of notation, in the next lemma with $J^+_{(\bar{A}_x, g')}$ it is denoted the set made of the diagonal of the compact $\bar{A}_x \times \bar{A}_x$ plus the pairs in $\bar{A}_x \times \bar{A}_x$ which can be joined by a continuous g' -causal curve of (M, g') entirely contained in $\bar{A}_x$ (it is an abuse of notation since $(\bar{A}_x, g')$ is not a spacetime as $\bar{A}_x$ is compact).

Lemma 3. *Let (M, g) be a compactly stably causal spacetime. Let C be a compact set and $B \supset C$ be a open set with compact closure. There is a metric $g_B \geq g$, $g_B > g$ on B , $g_B = g$ on $M \setminus B$, such that (M, g_B) at every point $x \in C$, admits an open neighborhood A_x with compact closure $\bar{A}_x \subset B$ such that $\bar{A}_x$ is g_B -causally convex. As a consequence, for every $g' \leq g_B$, $\bar{A}_x$ is g' -causally convex, no future inextendible continuous g' -causal curve is future imprisoned in $\bar{A}_x$, and $J^+_{(\bar{A}_x, g')}$ is compact.*

Proof. This proof is similar to that of [8, Lemma 3.10]. Since (M, g) is compactly stably causal there is g'_B , $g'_B > g$ on B , $g'_B = g$ on $M \setminus B$, such that (M, g'_B) is causal [9]. Let $g_B \geq g$ be a metric such that $g < g_B < g'_B$ on B , $g_B = g$ on $M \setminus B$. Let $x \in C$; it admits a nested family of g_B -globally hyperbolic neighborhoods V_n , $\bar{V}_{n+1} \subset V_n$, whose closures are all g_B -causally convex in V_1 , the set $\{V_n\}$ giving a base for the topology at x (see [13]). We can also assume that for all

n , $\bar{V}_n \subset B$, and V_1 has compact closure. If none of the sets $\bar{V}_n$ is g_B -causally convex in M there is a sequence of g_B -causal curves σ_n of endpoints x_n, z_n , with $x_n \rightarrow x$, $z_n \rightarrow x$, not entirely contained in V_1 and hence in $\bar{V}_2$. Let $c_n \in \bar{V}_2$ be the first point at which σ_n escapes $\bar{V}_2$, and let d_n be the last point at which σ_n reenters $\bar{V}_2$. Since $\bar{V}_2$ is compact there are $c, d \in \bar{V}_2$, and a subsequence σ_k such that $c_k \rightarrow c$, $d_k \rightarrow d$ and since V_1 is globally hyperbolic the causal relation on $V_1 \times V_1$, $J_{(V_1, g_B)}^+$, is closed and hence $(x, c), (d, x) \in J_{(V_1, g_B)}^+$ thus $d \neq c$ as the spacetime (V_1, g_B) is causal, finally $(x, c), (d, x) \in J_{(M, g_B)}^+$. Taking into account that $(c_k, d_k) \in J_{(M, g_B)}^+$ it is $(c, d) \in \bar{J}_{(M, g_B)}^+$.

Let us widen the light cones from g_B to g'_B . There is a g'_B -timelike curve connecting d to c passing through x , and since $(c, d) \in \bar{J}_{(M, g'_B)}^+$ and $I_{(M, g'_B)}^+$ is open there is a closed g'_B -timelike curve passing through x a contradiction with the causality of (M, g'_B) . The contradiction proves that there is a choice of n for which $\bar{V}_n$ is g_B -causally convex. Set $A_x = V_n$, then $\bar{A}_x$ is also clearly g' -causally convex for every $g' \leq g_B$. Since (V_1, g_B) is globally hyperbolic it is also non-total imprisoning, in particular no future inextendible continuous g' -causal curve is future imprisoned in the compact $\bar{A}_x$. The fact that $J_{(\bar{A}_x, g')}^+$ is compact follows from the compactness of $\bar{A}_x$, indeed by the limit curve theorem any sequence of continuous g' -causal curves in $\bar{A}_x$ with endpoints converging to a pair $(y, z) \in \bar{A}_x \times \bar{A}_x$, $y \neq z$, necessarily admits a limit g' -causal curve connecting y to z contained in $\bar{A}_x$, as the alternative would imply the presence of a future inextendible continuous g' -causal curve future imprisoned in the compact $\bar{A}_x$ passing through y .

Recall that if R^+ is a generic relation, $(R^+)^0$ is by definition the diagonal of $M \times M$, while $(R^+)^i$ denotes the composition of the relation with itself for i -times.

Lemma 4. *Let (M, g) , C , B , g_B and the sets $\{A_x\}$ be as in lemma 3. Let g' be a metric such that $g \leq g' \leq g_B$. Let $x \in C$, if (M, g') is K -causal then there is g'' , $g' \leq g'' \leq g_B$, such that (M, g'') is K -causal and $g'' > g$ on A_x .*

Proof. By assumption $K_{(M, g')}^+$ is antisymmetric.

Let $\tilde{g}$ be a metric such that $g' \leq \tilde{g} \leq g_B$, $\tilde{g} > g$ on A_x , $\tilde{g} = g'$ on $M \setminus A_x$ (e.g. a point dependent convex combination of g' and g_B).

For every $i \geq 0$, $K_{(M, g')}^+ \circ (J_{(\bar{A}_x, \tilde{g})}^+ \circ K_{(M, g')}^+)^i \subset K_{(M, \tilde{g})}^+$ as it is $J_{(\bar{A}_x, \tilde{g})}^+ \subset J_{(M, \tilde{g})}^+ \subset K_{(M, \tilde{g})}^+$ and $K_{(M, g')}^+ \subset K_{(M, \tilde{g})}^+$ (note that $K_{(M, \tilde{g})}^+$ is closed, transitive and contains $J_{(M, g')}^+$), thus

$$\bigcup_{i=0}^{+\infty} K_{(M, g')}^+ \circ (J_{(\bar{A}_x, \tilde{g})}^+ \circ K_{(M, g')}^+)^i \subset K_{(M, \tilde{g})}^+. \quad (1)$$

Suppose we prove that $\tilde{g}$ is also such that there is $N > 0$ so that

$$\bigcup_{i=0}^{+\infty} K_{(M, g')}^+ \circ (J_{(\bar{A}_x, \tilde{g})}^+ \circ K_{(M, g')}^+)^i = \bigcup_{i=0}^N K_{(M, g')}^+ \circ (J_{(\bar{A}_x, \tilde{g})}^+ \circ K_{(M, g')}^+)^i. \quad (2)$$

Each term $K_{(M,g')}^+ \circ (J_{(\bar{A}_x, \tilde{g})}^+ \circ K_{(M,g')}^+)^i$ is closed, a fact which follows easily from the observation that the composition of a closed a compact and a closed relation is closed. Thus the right-hand side of Eq. (2) is closed as it is the union of a finite number of closed sets. Moreover, it is also transitive because it equals the left-hand side of Eq. (2) which is clearly transitive. Finally, $J_{(M, \tilde{g})}^+$ is contained in it, a property which follows from the fact that since $\bar{A}_x$ is g' -causally convex it holds (g' and $\tilde{g}$ coincide outside A_x)

$$J_{(M, \tilde{g})}^+ = J_{(M, g')}^+ \cup J_{(M, g')}^+ \circ J_{(\bar{A}_x, \tilde{g})}^+ \circ J_{(M, g')}^+.$$

[The previous equation means that if $(x, z) \in J_{(M, \tilde{g})}^+$ then the $\tilde{g}$ -causal curve connecting x to z either passes outside A_x in which case it is g' -causal and $(x, z) \in J_{(M, g')}^+$ or it intersects A_x on, by g' -causal convexity of $\bar{A}_x$, a single segment. In this last case since the points at which the curve enters and escape $\bar{A}_x$ are $\tilde{g}$ -causal related, it is $(x, z) \in J_{(M, g')}^+ \circ J_{(\bar{A}_x, \tilde{g})}^+ \circ J_{(M, g')}^+$.]

Thus Eq. (2) implies

$$K_{(M, \tilde{g})}^+ \subset \bigcup_{i=0}^N K_{(M, g')}^+ \circ (J_{(\bar{A}_x, \tilde{g})}^+ \circ K_{(M, g')}^+)^i,$$

and hence by Eq. (1)

$$K_{(M, \tilde{g})}^+ = \bigcup_{i=0}^N K_{(M, g')}^+ \circ (J_{(\bar{A}_x, \tilde{g})}^+ \circ K_{(M, g')}^+)^i.$$

Consider a sequence of metrics $g'_n, g'_n \rightarrow g'$ pointwisely, which have the properties $g' \leq g'_n \leq g_B$, $g'_n > g$ on A_x , $g'_n = g'$ on $M \setminus A_x$ (for instance take $\bar{g}$, $g' \leq \bar{g} \leq g_B$, $\bar{g} > g$ on A_x , $\bar{g} = g'$ on $M \setminus A_x$ and define $g'_n = (1 - \frac{1}{n})g' + \frac{1}{n}\bar{g}$). Assume that a subsequence g'_k exists such that for each value of k , Eq. (2) with $\tilde{g} = g'_k$ does not hold no matter the value of $N(k)$. For every k since the equation

$$\bigcup_{i=0}^{+\infty} K_{(M, g')}^+ \circ (J_{(\bar{A}_x, g'_k)}^+ \circ K_{(M, g')}^+)^i = \bigcup_{i=0}^N K_{(M, g')}^+ \circ (J_{(\bar{A}_x, g'_k)}^+ \circ K_{(M, g')}^+)^i. \quad (3)$$

does not hold for any N , it is possible to find for each k a chain $x_j^{(k)}$ such that $(x_{2i}^{(k)}, x_{2i+1}^{(k)}) \in J_{(\bar{A}_x, g'_k)}^+$ and $(x_{2i+1}^{(k)}, x_{2i+2}^{(k)}) \in K_{(M, g')}^+ \setminus J_{(\bar{A}_x, g'_k)}^+$ with $0 \leq i \leq k$. Let D be an open set with compact closure such that $\bar{B} \subset D$. Note that for $1 \leq i \leq k-1$, since $J_{(\bar{A}_x, g')}^+ \subset J_{(\bar{A}_x, g'_k)}^+$

$$\begin{aligned} (x_{2i+1}^{(k)}, x_{2i+2}^{(k)}) &\in K_{(M, g')}^+ \cap \{\bar{A}_x \times \bar{A}_x\} \setminus J_{(\bar{A}_x, g')}^+ \\ &= [K_{(M, g')}^+ \cap (\bar{A}_x \times \bar{A}_x)] \setminus [J_{(M, g')}^+ \cap (\bar{A}_x \times \bar{A}_x)] \\ &= [K_{(M, g')}^+ \setminus J_{(M, g')}^+] \cap (\bar{A}_x \times \bar{A}_x) \end{aligned}$$

where the first equality follows from the g' -causal convexity of $\bar{A}_x$. Thus by lemma 1, there is $w_i^{(k)} \in \dot{D}$ such that $(x_{2i+1}^{(k)}, w_i^{(k)}) \in K_{(M, g')}^+$ and $(w_i^{(k)}, x_{2i+2}^{(k)}) \in K_{(M, g')}^+$.

Now we consider, by starting from $i = 1$, the sequence $(x_{2i}^{(k)}, x_{2i+1}^{(k)})$ as dependent on k and pass to a convergent subsequence $(x_{2i}^{(k_1)}, x_{2i+1}^{(k_1)})$, then we consider the sequence $(x_{2i+1}^{(k_1)}, w_i^{(k_1)})$ as dependent on k_1 and pass to a convergent subsequence $(x_{2i+1}^{(k_2)}, w_i^{(k_2)})$, then we consider the sequence $(w_i^{(k_2)}, x_{2i+2}^{(k_2)})$ as dependent on k_2 and pass to a convergent subsequence $(w_i^{(k_3)}, x_{2i+2}^{(k_3)})$, then we pass to the next value of i and continue in this way each time passing to a convergent subsequence.

Moreover, we use the fact that if an arbitrary sequence $(x_t^{(j)}, x_{t+1}^{(j)}) \in J_{(\bar{A}_x, g'_j)}^+$ converges to (x_t, x_{t+1}) then $(x_t, x_{t+1}) \in J_{(\bar{A}_x, g')}^+$ because $g'_j \rightarrow g'$ [recall that by the limit curve theorem since no g' -causal curve is future imprisoned in $\bar{A}_x$, a sequence of connecting g'_j -causal curves contained in $\bar{A}_x$ of endpoints $(x_t^{(j)}, x_{t+1}^{(j)})$ has a limit g' -causal curve contained in $\bar{A}_x$ of endpoints (x_t, x_{t+1})]. The limit pairs belong alternatively to $J_{(\bar{A}_x, g')}^+$, $K_{(M, g')}^+ \cap (\bar{A}_x \times \dot{D})$ and $K_{(M, g')}^+ \cap (\dot{D} \times \bar{A}_x)$, and since $J_{(\bar{A}_x, g')}^+ \subset K_{(M, g')}^+$ it is possible to find a sequence denoted y_s , $(y_s, y_{s+1}) \in K_{(M, g')}^+$, such that for s even, $y_s \in \dot{D}$, while for odd s , $y_s \in \bar{A}_x$. From this sequence we are going to find two points $p \in \bar{A}_x$ and $q \in \dot{D}$, and hence $p \neq q$, such that $(p, q) \in K_{(M, g')}^+$ and $(q, p) \in K_{(M, g')}^+$ in contradiction with $K_{(M, g')}^+$ -causality. Consider the sequence $(y_{2j+1}, y_{2j+2}) \in \bar{A}_x \times \dot{D}$ and pass to a converging subsequence $(y_{2j_r+1}, y_{2j_r+2}) \rightarrow (p, q)$. Since $K_{(M, g')}^+$ is closed, $(p, q) \in K_{(M, g')}^+$. Since $j_{r+1} \geq j_r + 1$, $2j_{r+1} + 1 \geq 2j_r + 2$ thus $(y_{2j_r+2}, y_{2j_{r+1}+1}) \in K_{(M, g')}^+$ as this last relation is transitive. Passing to the limit $r \rightarrow +\infty$, $(q, p) \in K_{(M, g')}^+$.

The contradiction proves that for sufficiently large n there is always $N(n)$ such that

$$\bigcup_{i=0}^{+\infty} K_{(M, g')}^+ \circ (J_{(\bar{A}_x, g'_n)}^+ \circ K_{(M, g')}^+)^i = \bigcup_{i=0}^{N(n)} K_{(M, g')}^+ \circ (J_{(\bar{A}_x, g'_n)}^+ \circ K_{(M, g')}^+)^i, \quad (4)$$

thus for sufficiently large n (in what follows we pass to a subsequence denoted in the same way so that it will hold for every n),

$$K_{(M, g'_n)}^+ = \bigcup_{i=0}^{N(n)} K_{(M, g')}^+ \circ (J_{(\bar{A}_x, g'_n)}^+ \circ K_{(M, g')}^+)^i.$$

We would conclude the proof by proving that there is a choice of n , such that the corresponding $K_{(M, g'_n)}^+$ is antisymmetric, indeed we would set $g'' = g'_n$.

Here the argument is basically the same that lead to the construction of points p and q . Since $K_{(M, g')}^+$ and $J_{(\bar{A}_x, g'_n)}^+$ are antisymmetric for every n , if $K_{(M, g'_n)}^+$ were not antisymmetric for no value of n then, for each n , we would find a closed chain of points so that the successive pairs belong to $J_{(\bar{A}_x, g'_n)}^+$ and $K_{(M, g')}^+ \setminus J_{(\bar{A}_x, g'_n)}^+$. However, a pair belonging to $K_{(M, g')}^+ \setminus J_{(\bar{A}_x, g'_n)}^+$ belongs also to $K_{(M, g')}^+ \setminus J_{(\bar{A}_x, g')}^+$ so that there is a point in $\dot{D}$ so as to split the pair in two, the middle point belonging to $\dot{D}$ and both pairs belonging to $K_{(M, g')}^+$. Then by

passing to subsequences as done above (basically to get the limit $n \rightarrow +\infty$), we find a chain of $K_{(M,g')}^+$ -related events alternatively belonging to $\bar{A}_x$ and $\bar{D}$. If the chain is finite and closed then it is easy to infer the contradiction that the spacetime is not $K_{(M,g')}^+$ -causal. If it is infinite one gets again the same conclusion by using the argument used above in the construction of p and q .

Lemma 5. *Let C be a compact. If (M, g) is K -causal then there is a metric $g_C \geq g$, such that $g_C > g$ on C , with (M, g_C) K -causal.*

Proof. Since (M, g) is K -causal it is compact stably causal. Let B , g_B and the sets $\{A_x\}$ be as in lemma 3. Since C is compact there is a finite covering $\{A_{x_i}\}$, thus one can start enlarging the metric in A_{x_1} while keeping K -causality according to lemma 4, and continue with successive enlargements so as to obtain a final metric g_C as in the statement of this lemma.

Theorem 1. *K -casuality coincides with stable causality.*

Proof. If (M, g) is K -causal then it is stably causal, indeed this result follows as a corollary of lemmas 2 and 5. The other direction is well known, see the discussion in the introduction.

Theorem 2. *If (M, g) is K -causal (stably causal) then $K^+ = J_S^+$.*

Proof. It is a consequence of theorem 6.2 of [8].

3. Causal easiness

The equivalence between K -causality and stable causality suggests to define a new conformal invariant property

Definition 1. *A spacetime which is A -causal and such that $A^+ = K^+$ is said to be causally easy.*

It is actually natural to define the property of causal easiness, indeed it appears in [9, Theorem 5] where it is proven that a spacetime which is chronological and has no lightlike line is causally easy. Notice that the condition $A^+ = K^+$ states that $\bar{J}^+$ is transitive.

Theorem 3. *(transverse conformal ladder) The compactness of the causal diamonds implies the closure of the causal relation which implies reflectivity which implies the transitivity of $\bar{J}^+$.*

Proof. It is well known that the compactness of the causal diamonds $J^+(x) \cap J^-(z)$ for all $x, z \in M$, implies $\bar{J}^+ = J^+$, see for instance [13, Prop. 3.68 and 3.71]. Now recall [12], that the relation $D^+ = \{(x, y) : y \in \overline{I^+(x)} \text{ and } x \in \overline{I^-(y)}\}$ is reflexive and transitive. It holds $D^+ = A^+$ iff the spacetime is reflective [12]. Since $J^+ \subset D^+ \subset A^+$, $J^+ = A^+$ implies reflectivity. Finally, reflectivity (but future or past reflectivity would be sufficient) implies the transitivity of A^+ , indeed $D^+ = A^+$ and since D^+ is transitive then A^+ is transitive.

A spacetime can be stably causal without being causally easy (the spacetime of figure 38 of [4] without the identification). A spacetime can also be causally easy without being causally continuous (1+1 Minkowski spacetime with a time-like geodesic segment removed). Causal easiness can indeed be placed between these two levels.

Theorem 4. *Causal continuity implies causal easiness which implies K -causality (stable causality).*

Proof. Recall [12] that a spacetime is causally continuous iff it is weakly distinguishing, that is D^+ is antisymmetric, and reflective, that is $D^+ = A^+$. But since D^+ is antisymmetric then A^+ is antisymmetric, that is the spacetime is A -casual. Moreover, by theorem 3 reflectivity implies the transitivity of A^+ ($A^+ = K^+$) thus the spacetime is causally easy.

Assume the spacetime is causally easy that is A^+ is antisymmetric and $A^+ = K^+$, then K^+ is antisymmetric, that is, the spacetime is K -causal.

The definition of causal easiness can be improved by weakening the condition of A -causality to strong causality.

Proposition 1. *A spacetime is causally easy iff it is strongly causal and $\bar{J}^+$ is transitive.*

Proof. To the right it is immediate since A -causality implies strong causality. Assume that the spacetime is strongly causal and $\bar{J}^+$ is transitive, and assume that the spacetime is not A -causal, then there are events x, z , $x \neq z$, such that $(x, z) \in \bar{A}^+$ and $(z, x) \in A^+$. Let σ_n be a sequence of causal curves of endpoints $(x_n, z_n) \rightarrow (x, z)$. By the limit curve theorem there is a limit causal curve σ^z ending at z (past inextendible or such that it connects x to z) and if $y \in \sigma^z \setminus \{z\}$ then $(x, y) \in \bar{J}^+$. Since $\bar{J}^+$ is transitive $(z, y) \in \bar{J}^+$ while clearly $(y, z) \in J^+$, thus by [7, theorem 3.4] the spacetime is not strongly causal, a contradiction.

In the definition of causal easiness the condition of causality cannot be further weakened to distinction, see figure 1.




Fig. 1. A distinguishing non-strongly causal, and hence non-causally easy, spacetime for which A^+ is transitive. Here the non-removed boundary at the bottom is identified with that at the top; as a consequence the spacetime is non-orientable but this feature is not essential. The only points at which strong causality is violated are those on the lightlike geodesic γ , and their future $A^+(x)$ is given by the shadowed region. This spacetime example is interesting because it shows that if strong causality is violated at x then there needs not to be a second event $z \neq x$, such that $x \in I^-(z) \cap \overline{I^+(z)}$.

Figure 2 summarizes the relationship between the various conformal invariant properties.




Fig. 2. The causal ladder of spacetimes with the levels involved in the implications which climb the ladder. Here the double arrow $A \Rightarrow B$ means that A implies B and that there are spacetime examples which show that properties A and B differ.

4. Conclusions

In this work the conjecture that K -causality and stable causality coincide has been proved. As a consequence in a K -causal spacetime the K^+ relation and the Seifert relation coincide. This is a powerful result which, once proved, allows to readily deduce several other results that otherwise should be obtained through more specific reasonings. Given this result it becomes also natural to introduce a new relation which I called causal easiness, which stays between causal continuity and stable causality.

I believe that the proof of the equivalence between stable causality and K -causality puts causality theory on a rather firm ground, especially for what concerns the levels from chronology up to stable causality.

References

1. Akolia, G. M., Joshi, P., and Vyas, U.: On almost causality. *J. Math. Phys.* **22**, 1243–1247 (1981)
2. Bernal, A. N. and Sánchez, M.: Globally hyperbolic spacetimes can be defined as ‘causal’ instead of ‘strongly causal’. *Class. Quantum Grav.* **24**, 745–749 (2007)
3. Dowker, H. F., Garcia, R. S., and Surya, S.: K -causality and degenerate spacetimes. *Class. Quantum Grav.* **17**, 4377–4396 (2000)
4. Hawking, S. W. and Ellis, G. F. R.: *The Large Scale Structure of Space-Time*. Cambridge: Cambridge University Press (1973)
5. Hawking, S. W. and Sachs, R. K.: Causally continuous spacetimes. *Commun. Math. Phys.* **35**, 287–296 (1974)
6. Janardhan, S. and Saraykar, R. V.: K -causal structure of space-time in general relativity. *Pramana-Journal of Physics* **70**, 587–601 (2008)
7. Minguzzi, E.: The causal ladder and the strength of K -causality. I. *Class. Quantum Grav.* **25**, 015009 (2008)
8. Minguzzi, E.: The causal ladder and the strength of K -causality. II. *Class. Quantum Grav.* **25**, 015010 (2008)
9. Minguzzi, E.: Chronological spacetimes without lightlike lines are stably causal (2008). Preprint: 0806.0153
10. Minguzzi, E.: Limit curve theorems in Lorentzian geometry (2008). *J. Math. Phys.* In press.

11. Minguzzi, E.: Non-imprisonment conditions on spacetime. *J. Math. Phys.* **49**, 062503 (2008)
12. Minguzzi, E.: Weak distinction and the optimal definition of causal continuity. *Class. Quantum Grav.* **25**, 075015 (2008)
13. Minguzzi, E. and Sánchez, M.: *The causal hierarchy of spacetimes*, Zurich: Eur. Math. Soc. Publ. House, vol. H. Baum, D. Alekseevsky (eds.), Recent developments in pseudo-Riemannian geometry, of *ESI Lect. Math. Phys.*, pages 299 – 358 (2008). gr-qc/0609119
14. Seifert, H.: The causal boundary of space-times. *Gen. Relativ. Gravit.* **1**, 247–259 (1971)
15. Senovilla, J. M. M.: Singularity theorems and their consequences. *Gen. Relativ. Gravit.* **30**, 701–848 (1998)
16. Sorkin, R. D. and Woolgar, E.: A causal order for spacetimes with C^0 Lorentzian metrics: proof of compactness of the space of causal curves. *Class. Quantum Grav.* **13**, 1971–1993 (1996)
17. Woodhouse, N. M. J.: The differentiable and causal structures of space-time. *J. Math. Phys.* **14**, 495–501 (1973)